

LATE-ONSET BEHAVIOURAL CHANGE: IS IT FRONTOTEMPORAL DEMENTIA?

Stefano F. Cappa

University Institute for Advanced Studies
(IUSS) and IRCCS Mondino Foundation,
Pavia, Italy

Disclosures

- Speaker, advisor: Roche, Biogen

Learning objectives

- Knowing about the multiple possible causes of late-onset behavioral changes in cognitive neurological practice
- Choosing the most appropriate strategy for differential diagnosis
- Planning the pharmacological and non-pharmacological management of the individual patient

Key message

Any clinically relevant behavioral change in an adult with a negative or incompatible history of psychiatric and neurological disorders needs to be comprehensively evaluated

References

- Borroni, B., Cosseddu, M., Pilotto, A., Premi, E., Archetti, S., Gasparotti, R., Cappa, S. and Padovani, A., 2015. Early stage of behavioral variant frontotemporal dementia: clinical and neuroimaging correlates. *Neurobiology of aging*, 36(11), pp.3108-3115.
- Ducharme, S., Pearl-Dowler, L., Gossink, F., McCarthy, J., Lai, J., Dickerson, B.C., Chertkow, H., Rapin, L., Vijverberg, E., Krudop, W. and Dols, A., 2019. The frontotemporal dementia versus primary psychiatric disorder (FTD versus PPD) checklist: a bedside clinical tool to identify behavioral variant FTD in patients with late-onset behavioral changes. *Journal of Alzheimer's Disease*, 67(1), pp.113-124.
- Ducharme, S., Dols, A., Laforce, R., Devenney, E., Kumfor, F., van den Stock, J., Dallaire-Théroux, C., Seelaar, H., Gossink, F., Vijverberg, E. and Huey, E., 2020. Recommendations to distinguish behavioural variant frontotemporal dementia from psychiatric disorders. *Brain*, 143(6), pp.1632-1650.